Computational Geometry-236719

(Spring 2019—Gill Barequet and Gil Ben-Shachar)

Assignment no. 3

Given: June 3, 2019 Due: **June 17, 2019**

Submission in **singletons**

Question 1.

Let L be a set of n lines in the plane. Give an $O(n \log n)$ -time algorithm to compute an axis-parallel rectangle that contains all the vertices of A(L) in its interior.

Question 2.

- 1. Let $S = \{p_1, ..., p_n\}$ (for $n \ge 3$) be the vertices of a regular convex polygon, and let C be its center. Let $P = S \cup C$. Prove that in the Voronoi diagram of P, the Voronoi cell of C contains n vertices. (A regular polygon is a polygon that is equiangular (all angles are equal in measure) and equilateral (all sides have the same length).
- 2. Assuming general position, prove that for a Voronoi diagram of n points, where n is large enough, the average number of vertices of a cell is 6.

Question 3.

GG(S), the *Gabriel Graph* of a point set S in the plane, is defined as follows: Two points $p, q \in S$ are connected by an edge of the graph if the circle with diameter pq does not contain any other point of S in its interior.

- 1. Prove that DT(S) (Delaunay Triangulation of S) contains the Gabriel graph of S.
- 2. Prove that p and q are adjacent in GG(S) iff the Delaunay edge that connects between them intersects its dual Voronoi edge.
- 3. Give an $O(n \log n)$ -time algorithm to compute the Gabriel graph of a set of n points.

Question 4.

Let S be a set of n points in the plane, and let t be the number of lines that pass through exactly \sqrt{n} points of S. Prove that $t = O(\sqrt{n})$.