

International Journal of Shape Modeling
Vol. 9, No. 2 (2003) 191–201
© World Scientific Publishing Company

Intrinsic Morphing of Compatible Triangulations

VITALY SURAZHSKY*
vitus@cs.technion.ac.il

CRAIG GOTSMAN
gotsman@cs.technion.ac.il

*Center for Graphics and Geometric Computing,
Department of Computer Science,
Technion—Israel Institute of Technology,
Haifa 32000, Israel*

Two planar triangulations with a correspondence between two vertex sets are compatible (*isomorphic*) if they are topologically equivalent. This work presents a simple and robust method for morphing two compatible planar triangulations with identical convex boundaries that locally preserves the intrinsic geometric properties of triangles throughout the morph. The method is based on the barycentric coordinates representation of planar triangulations, and thus, guarantees compatibility of all intermediate triangulations. The intrinsic properties are preserved by interpolating angles and edge lengths components of *mean value* barycentric coordinates, rather than interpolating the barycentric coordinates themselves. As a result, the method generates a natural-looking and guaranteed intersection-free morphing sequence.

Keywords: Controllable morphing; Compatible triangulations; Morphing; Self-intersection elimination

1. Introduction

Morphing, also known as metamorphosis, is the gradual and continuous transformation of one shape (the *source*) into another (the *target*). Morphing has wide practical use in areas such as computer graphics, animation and modeling. To achieve more spectacular, impressive and accurate results, the morphing process requires a lot of the work to be done manually. A major research challenge is to develop techniques that will automate this process as much as possible.

The morphing problem has been investigated in many contexts, e.g., morphing of two-dimensional images^{1,2,3}, polygons and polylines^{4,5,6,7,8,9,3}, freeform curves¹⁰ and even voxel-based volumetric representations.¹¹ The morphing process always consists of solving two main problems. The first one is to find a correspondence

*Currently at Department of Informatics, Oslo University, Oslo, Norway

between elements (features) of the two shapes. The second problem is to find trajectories that corresponding elements traverse during the morphing process. Regrettably, a formal definition of a successful correspondence does not exist, as well as a definition of a successful solution of the trajectory problem. In this work we assume the more typical morphing scenario, namely, that the correspondence is given and only the trajectory problem is to be solved.

The naive approach to solve the trajectory problem is to choose the trajectories to be straight lines, where every feature of the shape travels with a constant velocity along its line towards the corresponding feature of the target during the morph. Unfortunately, this simple approach can lead to undesirable results. The intermediate shapes can vanish, i.e. degenerate into a single point, or contain self-intersections even though the source and target are simple, namely, self-intersection-free. Even if the linear morph is free of self-intersections and degeneracies, its intermediate shapes may have areas or distances between features far from those of the source and target, resulting in a “misbehaved” looking morph. Most of the research on solving the trajectory problem for morphing concentrates on trying to eliminate self-intersections and preserve the geometrical properties of the intermediate shapes. Many existing methods achieve good results for many inputs, however, none is able to guarantee any properties of the resulting morph.

Triangulations are ubiquitously used in computer graphics as a representation and a parameterization (e.g., for texture mapping) of surfaces and planar shapes. Two triangulations with a correspondence between their vertex sets are said to be compatible (*isomorphic*), if they are topologically equivalent. Floater and Gotsman¹² introduced an innovative approach for morphing compatible planar triangulations. This approach is based on the barycentric coordinates representation of planar triangulations. The morphing interpolation procedure is done in the space of barycentric coordinates, rather than in vertex geometry space. This approach is the only one, which applied to compatible triangulations with identical boundaries, is able to analytically guarantee a continuous sequence of valid intermediate triangulations, namely, the triangulation does not self-intersect during the morph.

Surazhsky and Gotsman¹³ analyzed the basic approach for morphing compatible triangulations,¹² investigated its properties and capabilities and presented several extensions that allow control over the properties of morphs. They showed that the basic approach has many degrees of freedom, which can be utilized in order to obtain morphs with various characteristics and still be free of self-intersections. However, the schemes to obtain more natural morphs proposed by Surazhsky and Gotsman¹³ are rather complicated and have problems inherent to the barycentric coordinates used in that work.

In this work we exploit the non-uniqueness of barycentric coordinates and use mean value barycentric coordinates that were recently introduced by Floater.¹⁴ These coordinates reflect the shapes of triangles in a way similar to coordinates

used in harmonic or conformal mappings.¹⁵ However, the naive linear interpolation of mean value coordinates cannot guarantee that the shapes of triangles within intermediate triangulations transform uniformly, and thus, naturally. Mean value coordinates depend solely on triangle angles and edge lengths within a triangulation. Hence, in order to achieve a uniform transformation of the triangles shapes we interpolate angles and edge lengths of the corresponding triangles, and then use the intermediate values to generate intermediate mean value coordinates.

Guaranteed intersection-free morphing of compatible planar triangulations allows morphing of any compatible piecewise linear objects such as polygons, poly-lines, tilings and so called *stick figures* in general. Formally, stick figures are defined as connected straight-line plane graphs. Guaranteed intersection-free morphing of stick figures is achieved by embedding them as a subset of edges within compatible triangulations with identical boundaries. This embedding technique was introduced³ for polygons and later extended¹⁶ for stick figures.

Constructing compatible triangulations for embedding stick figures is a difficult problem and most of the existing algorithms^{17,18,19,16} produce triangulations that may have long, skinny and close to degenerate triangles together with many additional (*Steiner*) vertices. Moreover, the resulting compatible triangulations must usually be refined in order to create more natural morphing sequences.⁴ The refinement increases the number of vertices by a factor of 10–100 on the average, and this leads to very high computational cost of morphing sequences that can easily reach hours of CPU time even on modern 2-3GHz computers. Surazhsky and Gotsman²⁰ presented a novel algorithm, which is able to construct compatible triangulations with a very small (close to optimal) number of Steiner vertices. However, having many of the triangles close to degenerate, the resulting compatible triangulation must be refined in order to be used with existing morphing algorithms.

The contribution of this paper is a relatively simple method to generate natural morph sequences between two planar compatible triangulations. It is also suitable for generating a natural morph between two stick figures. Our method is robust because the quality of the embedding triangulations used to morph two stick figures does not significantly affect the generated morphing sequence. Thus, our method can be applied to compatible triangulations with a very small number of Steiner vertices to obtain natural morphs in real time.

The paper is organized as follows: Section 2 introduces the notions of compatible triangulations and barycentric coordinates and define barycentric coordinates representation for triangulations. Section 3 shows how to produce a self-intersection free morph between compatible triangulations and then extends it to generate the intrinsic morph. Experimental results generated by our algorithm are shown in Section 4, and we conclude in Section 5.




Fig. 1. (a) Triangulation of 9 points in the plane. (b) Triangulation not compatible with (a). Vertex correspondence coded in digits. (c) Triangulation compatible with (a).

2. Background

A simple graph $G = G(V, E)$ is a set of vertices $V = \{1, \dots, |V|\}$ and a set of edges E , such that E is a subset of all unordered pairs of vertices $\{i, j\}$, when $i \neq j$. Two graphs G_0 and G_1 are *isomorphic* if there is a 1–1 correspondence between their vertices and edges in such a way that corresponding edges link corresponding vertices. A simple graph G is said to be *planar* if it can be drawn in the plane using simple curves for edges and the only intersections between the curves are at common endpoints standing for vertices. This representation of a planar graph G is called a *plane graph*, denoted by an ordered pair (G, \wp) . A mapping $\wp: V \rightarrow \mathbb{R}^2$ is a point sequence $\{i \mapsto (x_i, y_i) \mid i: 1, \dots, |V|\}$. The following notations are equivalent and denote the coordinates of vertex i : $\wp(i)$, p_i , (x_i, y_i) .

A plane graph partitions the plane into connected regions called *faces*. Obviously, a plane graph has a single unbounded face, denoted by the *outer* face. A plane graph is said to be *triangulated* if all its bounded faces have exactly three edges. A *planar triangulation* $\mathcal{T} = \mathcal{T}(G, \wp)$ is a simple triangulated plane graph such that its edges are represented by straight lines. We call a triangulation *valid* if it satisfies the above definition, in particular, the only intersections between its edges are at common endpoints.

Two (valid) triangulations are *isomorphic* if their planar graphs are isomorphic and the corresponding faces have the same orientation. To be consistent with other works, we also call isomorphic triangulations *compatible* triangulations. In this paper we will deal with compatible triangulations whose corresponding vertices have the same indices, namely, their planar graphs are identical. Thus, if $\mathcal{T}_0 = \mathcal{T}(G_0, \wp_0)$ and $\mathcal{T}_1 = \mathcal{T}(G_1, \wp_1)$ are compatible then $G_0 = G_1$. Fig. 1 shows some compatible and non-compatible triangulations.

2.1. Barycentric Coordinates

Given a polygon with k vertices p_1, p_2, \dots, p_k , $k \geq 3$, any point p in the plane can be expressed as:

$$p = \sum_{i=1}^k \lambda_i \cdot p_i, \quad \sum_{i=1}^k \lambda_i = 1. \quad (1)$$




Fig. 2. Elements of mean value coordinates.

The coefficients $\lambda_1, \dots, \lambda_k$ in these equations are said to be *barycentric coordinates* of p relative to p_1, \dots, p_k . When p lies in the convex hull of a polygon, it can be expressed as a convex combination of the polygon vertices, namely, all barycentric coordinates have positive values. In this paper we consider the case when p lies in the kernel of a star-shaped polygon.

In the special case when the polygon has three vertices, namely, p lies in a triangle $\triangle(p_1, p_2, p_3)$, the barycentric coordinates of p with respect to p_1, p_2, p_3 are uniquely determined by (1). Barycentric coordinates of p with respect to polygons with $k > 3$ vertices are not unique. A simple solution is to choose any triangle containing p whose vertices are vertices of the polygon. Barycentric coordinates of p may now be defined as non-zero coordinates with respect to the triangle vertices, and zeros for all other vertices. However, it has long been a challenging problem to find *strictly positive* barycentric coordinates that *continuously* and *smoothly* depend on p and the polygon vertices. Methods proposed by various works^{21,22,13,15} failed to satisfy at least one of the above requirements, and only recently Floater¹⁴ solved this problem. He introduced the following simple formula for barycentric coordinates:

$$\lambda_i = \frac{w_i}{\sum_{j=1}^k w_j}, \quad w_i = \frac{\tan\left(\frac{\alpha_i}{2}\right) + \tan\left(\frac{\alpha_{i-1}}{2}\right)}{\ell_i}. \quad (2)$$

α_i is the angle at p of the triangle $\triangle(p, p_i, p_{i+1})$ and ℓ_i is the length of edge (p, p_i) , namely, $\ell_i = \|p - p_i\|$. See Fig. 2(a). Since $0 < \alpha_i < \pi$ being a triangle angle, it is easy to see that resulting barycentric coordinates are strictly positive. The barycentric coordinates defined in (2) are called *mean value* coordinates inspired by the Mean Value theorem for harmonic functions. Mean value coordinates approximate harmonic maps by piecewise linear maps over triangulations. This important capability of mean value coordinates to preserve angles is exploited by our morphing scheme to produce natural morphs.

2.2. Drawing Triangulated Graphs

It has been shown by Fáry²³ that every planar graph has a *straight line representation*. Hence, for every planar triangulated graph G , there exists a point sequence

\wp such that $\mathcal{T} = \mathcal{T}(G, \wp)$ is a (valid) triangulation. Tutte²⁴ described the following method to generate \wp : The boundary vertices of G are mapped to an arbitrary convex polygon with the same number of vertices and the same vertex order. Then, the interior vertices are placed such that every vertex is the centroid of its neighboring vertices. This scheme was extended by Floater.²¹ Each interior vertex can be *any* convex combination of its neighbors. In terms of barycentric coordinates, any positive barycentric coordinates for each interior vertex may be chosen with respect to its neighbors.

To compute \wp , we use the following method. Let $G = G(V, E)$ be a simple triangulated graph, with $|V| = N$. We assume that boundary vertices of G have been identified. Let V_I be the set of the interior vertices and V_B be the set of the boundary vertices such that $|V_I| = n$ and $|V_B| = N - n = k$. Without loss of generality assume $V_I = \{1, \dots, n\}$ and $V_B = \{n+1, \dots, N\}$.

Now we wish to find coordinates of the graph vertices, namely, for each vertex $i \in V$ to find $\wp(i) = p_i = (x_i, y_i)$. We define \wp for each vertex $i \in V_B$ to be coordinates of the vertices of a k -sided convex polygon.

For each interior vertex we may choose arbitrary non-negative barycentric coordinates relative to its neighbors, namely, for each vertex $i \in V_I$ a set of scalars $\lambda_{i,j}$ for $j = 1, \dots, N$ such that

$$\begin{aligned} \lambda_{i,j} &= 0, \quad \{i, j\} \notin E, \\ \lambda_{i,j} &\geq 0, \quad \{i, j\} \in E, \end{aligned} \quad \begin{array}{l} \text{barycentric coordinate} \\ \text{of } i \text{ relatively to } j. \end{array} \quad (3)$$

Next we define p_1, \dots, p_n to be the solution of the following system of linear equations

$$p_i = \sum_{j=1}^N \lambda_{i,j} p_j, \quad i = 1, \dots, n. \quad (4)$$

This system contains n linear equations with n variables. It was shown by Floater²¹ that the corresponding matrix of these equations is non-singular. Therefore, a unique solution always exists. Thus, barycentric coordinates defined in (1) together with the locations of the triangulation boundary vertices uniquely determine the locations of the interior vertices. This representation is called *barycentric coordinates representation* of the triangulation.

Bi-conjugate gradient methods can be efficiently used to numerically solve the non-symmetric linear system (4). For even more efficient multilevel solvers see the recent work of Aksoylu et al.²⁵

3. Morphing Triangulations

A *morph* between two compatible triangulations $\mathcal{T}_0 = \mathcal{T}(G, \wp_0)$ and $\mathcal{T}_1 = \mathcal{T}(G, \wp_1)$ is a gradual transformation of \mathcal{T}_0 into \mathcal{T}_1 . This transformation may be viewed as a continuous function $\mathcal{T}(t)$, $t \in [0, 1]$ and $\mathcal{T}(0) = \mathcal{T}_0$, $\mathcal{T}(1) = \mathcal{T}_1$. A morph between \mathcal{T}_0

and \mathcal{T}_1 is *valid* if for all $t \in [0, 1]$, $\mathcal{T}(t)$ is a valid triangulation compatible with \mathcal{T}_0 (and \mathcal{T}_1). In this work we consider triangulations \mathcal{T}_0 and \mathcal{T}_1 such that the boundaries of the triangulations coincide. The boundaries of $\mathcal{T}(t)$ for $0 \leq t \leq 1$ will naturally also coincide, namely, $p_i(t) = p_i(0) = p_i(1)$ for $n < i \leq N$ and $0 \leq t \leq 1$. To find $\mathcal{T}(t)$ means to find $p_i(t)$ for $1 \leq i \leq n$ such that $\mathcal{T}(G, \wp_t)$ is compatible with \mathcal{T}_0 .

Floater and Gotsman¹² proposed to interpolate barycentric coordinates of the interior vertices of \mathcal{T}_0 and \mathcal{T}_1 rather than the interior vertex locations. Barycentric coordinates of \mathcal{T}_0 and \mathcal{T}_1 can be any valid positive barycentric coordinates, but strictly positive mean value coordinates defined in (2) are currently the best choice due to their properties. The linear interpolation of strictly positive barycentric coordinates yields strictly positive intermediate barycentric coordinates as well. The intermediate barycentric coordinates uniquely define valid intermediate triangulations, resulting in a valid morph. Formally, we denote barycentric coordinates defined in (3) for \mathcal{T}_0 and \mathcal{T}_1 as $\lambda_{i,j}(0)$ and $\lambda_{i,j}(1)$, respectively. Intermediate barycentric coordinates are denoted by $\lambda_{i,j}(t)$ for $0 < t < 1$ and defined as the simple linear interpolations:

$$\lambda_{i,j}(t) = (1 - t) \cdot \lambda_{i,j}(0) + t \cdot \lambda_{i,j}(1). \quad (5)$$

Note that we need to solve the linear system (4) for every t of the morphing sequence. As an initial guess, one could use the solution of the previous t in the morphing sequence.

3.1. Intrinsic Morphing

In this work we go one step further and linearly interpolate components of the mean value coordinates formula. In other words, we interpolate the triangle angles and the edge lengths, rather than just the mean value coordinates themselves.

Formally, instead of (5) we use:

$$\begin{aligned} \alpha_{i,j}(t) &= (1 - t) \cdot \alpha_{i,j}(0) + t \cdot \alpha_{i,j}(1) \\ \ell_{i,j}(t) &= (1 - t) \cdot \ell_{i,j}(0) + t \cdot \ell_{i,j}(1) \end{aligned} \quad (6)$$

and substitute them into (2) to obtain $\lambda_{i,j}(t)$. $\ell_{i,j}$ is the length of edge (i, j) and $\alpha_{i,j}$ is the angle at vertex i adjacent from the left to edge $\{i, j\}$. See Fig. 2(b). $\alpha_{i,j}(0)$, $\alpha_{i,j}(1)$ and $\ell_{i,j}(0)$, $\ell_{i,j}(1)$ for all (i, j) such that $\{i, j\} \in E$ are taken from \mathcal{T}_0 and \mathcal{T}_1 , respectively.

The motivation behind this was an attempt to obtain as uniform as possible transformations of intrinsic properties of triangulations throughout the morphing process. Since the mean value coordinates formula (2) is not linear in the angles and the edge lengths, the linear interpolation of mean value coordinates corresponds to some strictly nonlinear transformations of the angles and the edge lengths. This may result in transformations of the triangle angles and the edge lengths that are significantly far from uniform, even though the intermediate triangulations are still valid.

Note that the linear interpolation of the angles and the edge lengths does not guarantee that all intermediate triangulations will have these exact angles and edge lengths. Usually, such triangulations simply do not exist. If they did exist, there would be no need to construct intermediate triangulations using mean value barycentric coordinates. One could construct intermediate triangulations directly using the angles and edge lengths. Thus, our morphing method, being based on the angle-preserving property of mean value coordinates, produces intermediate triangulations that approximate as close as possible linearly interpolated angles. This approximation is similar to the local angle-preserving properties of conformal mapping of 3D triangulations, which aims to preserve the angle between any two edges with a common endpoint. Note that having this angle-preserving property, our method guarantees, at the same time, that all intermediate triangulations are still valid.

It is important to emphasize the principal factor that makes it possible to obtain guaranteed intersection-free morphing of compatible triangulations and then to extend it to intrinsic morphing. The key lies in changing representation of the triangulation such that the space where we perform the linear interpolation (to generate a morphing sequence) is extended. This means that we interpolate more elements in order to obtain better morphs. The straightforward representation of the triangulation geometry, namely, the vertex locations gives $2n$ (x 's and y 's) degrees of freedom. The barycentric coordinates representation involves $\sum_{i=1}^n d_i > 3n$ elements ($\lambda_{i,j}$'s, $\approx 6n$ for large triangulations), where d_i is the degree of vertex i . The intrinsic morph interpolates $\sum_{i=1}^n d_i$ angles ($\alpha_{i,j}$'s) and $|E| - k$ edges ($\ell_{i,j}$'s), in total approximately $9n$ elements for large triangulations. Thus, every time the interpolation space is enlarged, morphs with better characteristics may be obtained.

4. Experimental Results

In practice, morphing is performed more frequently on stick figures than on planar triangulations. Thus, the problem of morphing planar figures is reduced to that of morphing triangulations, and the edges that are not part of the figure are ignored in the resulting morph. Figures 3 and 4 show morphs between two polylines, the shapes of the two letters U and S , and two stick figures, the shapes of a scorpion and a dragonfly. These examples have been embedded within compatible triangulations using techniques developed in ²⁰. Note that the triangulations in Fig. 3(a)–(b) and Fig. 4(a)–(b) have many long and skinny triangles, since the number of Steiner vertices is close to minimal. On the other hand, the triangulations in Fig. 3(c)–(d) do not contain long and skinny triangles, but have many Steiner vertices. The examples show that applying interpolation of barycentric coordinates on triangulations with long, skinny triangles generates strange-looking morphs, see Fig. 3(f),(g) and Fig. 4(d). Using the same technique on high quality triangulations produces rather natural morphs, see Fig. 3(h),(i). However, while the given polylines can be viewed as an approximation of smooth curves, the intermediate polylines are no longer

smooth, no matter what barycentric coordinates were used. On the other hand, the intrinsic morph produces natural morphs for high quality triangulations—see Fig. 3(j)—as well as for triangulations with long and skinny triangles, see Fig. 3(k). Moreover, for both these morphs the polylines stay smooth throughout the morphing sequences due to the angle-preserving property of our method. This shows that the intrinsic morphing scheme is robust with respect to the triangulations used to perform morphing.

5. Discussion and Conclusion

We have presented a simple, robust and efficient method for morphing compatible triangulations that can be used to produce very natural and guaranteed intersection-free morphing sequences between stick figures. The quality and behavior of the resulting morphs is not affected by the quality of the embedding triangulations. Even triangulations containing many triangles which are close to degenerate may be used to obtain natural morphs. This allows the use of compatible triangulations with a very small number of Steiner vertices, obtaining intersection-free natural morphs in almost real time.

References

1. T. Beier and S. Neely, “Feature-based image metamorphosis,” *Computer Graphics (SIGGRAPH '92)*, vol. 26, no. 2, pp. 35–42, 1992.
2. A. Tal and G. Elber, “Image morphing with feature preserving texture,” *Computer Graphics Forum (Eurographics '99 Proceedings)*, vol. 18, no. 3, pp. 339–348, 1999.
3. C. Gotsman and V. Surazhsky, “Guaranteed intersection-free polygon morphing,” *Computers and Graphics*, vol. 25, no. 1, pp. 67–75, 2001.
4. M. Alexa, D. Cohen-Or, and D. Levin, “As-rigid-as-possible polygon morphing,” *Proceedings of SIGGRAPH '2000*, pp. 157–164, 2000.
5. E. Carmel and D. Cohen-Or, “Warp-guided object-space morphing,” *The Visual Computer*, vol. 13, pp. 465–478, 1997.
6. E. Goldstein and C. Gotsman, “Polygon morphing using a multiresolution representation,” *Proceeding of Graphics Interface*, pp. 246–254, 1995.
7. T. W. Sederberg, P. Gao, G. Wang, and H. Mu, “2D shape blending: an intrinsic solution to the vertex path problem,” *Computer Graphics (SIGGRAPH '93)*, vol. 27, pp. 15–18, 1993.
8. T. W. Sederberg and E. Greenwood, “A physically based approach to 2D shape blending,” *Computer Graphics (SIGGRAPH '92)*, vol. 26, pp. 25–34, 1992.
9. M. Shapira and A. Rappoport, “Shape blending using the star-skeleton representation,” *IEEE Trans. on Computer Graphics and Application*, vol. 15, no. 2, pp. 44–51, 1995.
10. T. Samoilov and G. Elber, “Self-intersection elimination in metamorphosis of two-dimensional curves,” *The Visual Computer*, vol. 14, pp. 415–428, 1998.
11. D. Cohen-Or, D. Levin, and A. Solomovici, “Three-dimensional distance field metamorphosis,” *ACM Transactions on Graphics*, vol. 17, pp. 116–141, Apr. 1998.
12. M. S. Floater and C. Gotsman, “How to morph tilings injectively,” *Computational and Applied Mathematics*, vol. 101, pp. 117–129, 1999.




Fig. 3. (a)–(b) Two polylines of letters U and S are embedded into compatible triangulations with 12 Steiner vertices. (c)–(d) The same polylines are embedded into high quality compatible triangulations with 186 Steiner vertices. (e) The linear interpolation of vertex positions results in an invalid morph. (f),(g) The linear interpolation of mean value (f) and minimal variance¹³ (g) barycentric coordinates results in a valid but unnatural morph whose intermediate polylines behave strangely. (h),(i) Linear interpolation of mean value (h) and minimal variance¹³ (i) coordinates applied to the high quality triangulations (c)–(d) produces a natural morph, but the intermediate polylines are no longer smooth. (j) The intrinsic morph of the high quality triangulations (c)–(d) behaves naturally and has smooth intermediate polylines. (k) The intrinsic morph between triangulations (a)–(b) is also natural, has smooth intermediate polylines and is very similar to (j), even though a very small number of Steiner vertices are used.

13. V. Surazhsky and C. Gotsman, “Controllable morphing of compatible planar triangulations,” *ACM Transactions on Graphics*, vol. 20, no. 4, pp. 203–231, 2001.
14. M. S. Floater, “Mean value coordinates,” *CAGD*, 2003. To appear.
15. M. Desbrun, M. Meyer, and P. Alliez, “Intrinsic parameterizations of surface meshes,” *Proceedings of EUROGRAPHICS 2002*, vol. 21, no. 2, 2002.
16. V. Surazhsky and C. Gotsman, “Morphing stick figures using optimized compatible triangulations,” *Proceedings of Pacific Graphics*, pp. 40–49, Oct. 2001.
17. B. Aronov, R. Seidel, and D. L. Souvaine, “On compatible triangulations of simple polygons,” *Computational Geometry: Theory and Applications*, vol. 3, pp. 27–35, 1993.
18. E. Kranakis and J. Urrutia, “Isomorphic triangulations with small number of Steiner points,” *International Journal of Computational Geometry & Applications*, vol. 9, no. 2, pp. 171–180, 1999.
19. H. Gupta and R. Wenger, “Constructing piecewise linear homeomorphisms of simple polygons,” *Journal of Algorithms*, vol. 22, no. 1, pp. 142–157, 1997.
20. V. Surazhsky and C. Gotsman, “High quality compatible triangulations,” *Proceedings of 11th International Meshing Roundtable*, pp. 183–192, Sept. 2002.
21. M. S. Floater, “Parameterization and smooth approximation of surface triangulation,”




Fig. 4. Morphing between figures of a scorpion and a dragonfly embedded into compatible triangulations: (a)–(b) Compatible embedding triangulations with only 11 Steiner vertices. (c) The linear interpolation of vertex positions results in an invalid morph. (d) The linear interpolation of mean value coordinates results in a valid but unnatural morph having a strange behavior of the tail components. (e) The intrinsic morph behaves naturally and the tail components evolve smoothly in the intermediate triangulations.

Computer Aided Geometric Design, vol. 14, pp. 231–250, 1997.

22. K. Sugihara, “Surface interpolation based on new local coordinates,” *Computer-Aided Design*, vol. 31, pp. 51–58, 1999.
23. I. Fáry, “On straight line representation of planar graphs,” *Acta Univ. Szeged Sect. Sci. Math.*, vol. 11, pp. 229–233, 1948.
24. W. T. Tutte, “How to draw a graph,” *Proc. London Math. Soc.*, vol. 13, pp. 743–768, 1963.
25. B. Aksoylu, A. Khodakovsky, and P. Schröder, “Multilevel solvers for unstructured surface meshes,” Submitted.